



***Lens culinaris* Medikus a traditional crop of Kumaun Himalaya and ethnobotanical prespectives**

Kamal Singh Rawat¹, Manju Lata Upadhyaya²

^{1,2} Department of Botany, Kumaun University, S.S.J. Campus Almora, Uttarakhand, India

Abstract

Lentils (*Lens culinaris*) is most nutritious annual cool season crop belongs to Fabaceae family commonly called Lentil also known by many common names viz., Masur (small seeded), (Kay, 1979). Lentil was one of the first domesticated grain legumes, originating from the Near East center of origin (Zohary, 1999). Lentil subsequently spread to central Asia and the Mediterranean Basin (Cubero, 1981; Lev-Yadun *et al.*, 2000) among grain legumes, lentil seeds have highest protein content (22% to 34.6%) and considered as the third highest level of protein of any legume after soybeans and hemp, and major source of protein in vegetarian diet. Its lens-shaped seeds exist with a spectrum of colors including yellow, red, orange, green, brown or black depending on the cultivar, and subsequently the composition of the seed coats and cotyledons (Xu and Chang, 2010).

Keywords: Kumaun, ethnobotanical, Himalaya, cultivar

Introduction

Lens culinaris is a much-branched, erect, bushy, annual plant growing 30 - 80cm tall. Lentil is one of the oldest pulse crops and is of ancient cultivation in western Asia, Egypt and southern Europe. It probably originated in western Asia, from where it eventually spread throughout temperate and subtropical regions of the world and into the higher elevations in tropical areas. The earliest records of it being used date back

6 to 11,000 BC in Ancient Greece, though it was another 5,000 years before it was being cultivated regularly. A very important food plant, its nutritious seed has become a staple food for many (Facciola, 1990) [6]. The species is hermaphrodite (has both male and female ofloral parts) and is pollinated by Cleistogomy (self-pollinating without flowers ever opening). The plant is self-fertile and useful for crop rotation since it has the nitrogen fixing ability.

Origin and Geographical Distribution

Lentil is one of the oldest pulse crops and of ancient cultivation in western Asia, Egypt and southern Europe. It probably originated in western Asia, from where it spread into the Mediterranean region, Asia, Africa and Europe. Lentil was a common part of the diet of the ancient Greeks, Jews and Romans and was the mainstay of the poor, especially in Egypt. It was associated with many legends, tales and customs, and it is the first pulse crop mentioned in the Bible. The oldest archaeological remains of lentil are from Greece, dated 11,000 BC, and Syria, dated 8500–7500 BC. However, it is uncertain whether they were from cultivated plants or from wild ones. India.

Botanical Discription

The botanical features of *Lens culinaris* (cultivated lentil) can be described as annual bushy herb, slender almost erect or suberect, much-branched, softly hairy; stems slender, angular, 15-75 cm height (Duke, 1981; Muehlbauer *et al.*, 1985) [5]. Ten to sixteen leaflets are subtended on the rachis (40-50 mm); upper leaves have simple tendrils while lower leaves are mucronate

(Muehlbauer *et al.*, 1985). "The leaves are alternate, compound, pinnate, usually ending in a tendril or bristly; leaflets 4-7 pairs(fig-2), alternate or opposite; oval, sessile, 1-2 cm long; stipules small, entire; stipules absent; pods oblong, flattened or compressed, smooth, to 1.3 cm long, 1-2-seeded; seed biconvex, rounded, small, 4-8 mm × 2.2-3 mm, lens-shaped, green, greenish-brown or light red speckled with black; the weight of 100 seeds range from 2 to 8 g; cotyledons red, orange, yellow, or green, bleaching to yellow, often showing through the testa, influencing its apparent color" (Kay, 1979; Duke, 1981; and Muehlbauer *et al.*, 1995) [5]. Flowers are small, pale blue, purple, white or pink, in axillary 1-4-flowered racemes; 1-4 flowers are borne on a single peduncle and a single plant can produce upto 10-150 peduncles each being 2.5-5 cm long (Muehlbauer *et al.*, 1985). Flowering proceeds acropetally. The size of seeds increase from the types grown in eastern regions to western types. Two types, namely; macrosperma, found mainly in the Mediterranean region and the New World (seed size ranging from 6 to 9 mm in diameter and yellow cotyledons with little or no pigmentation), and microsperma (2 to 6 mm with red orange or yellow cotyledons) found on the Indian subcontinent(fig-1) (Hawtin *et al.*, 1980; Muehlbauer *et al.*, 1985) [4]. The first one includes the Chilean or yellow cotyledon types while the latter includes the small seeded Persian or red cotyledon lentils (Kay, 1979). Germination is hypogeal and this keeps the developing seedlings below ground level which reduces the effects of freezing and other desiccating environmental conditions (Muehlbauer *et al.*, 1985).

Climate

Seeds will germinate at temperatures above freezing but best at the range of 18-21°C; temperatures above 27°C are harmful; optimum temperatures for growth and yields are around 24°C. Lentils are grown as a cool weather or winter crop in the semi-arid tropics, cultivated from sea level to 3,800 m, but are not

suited to the humid tropics. They are less damaged by drought than by waterlogging. Lentils thrive well on a wide range of soils from light loams and alluvial soil to black cotton soils, best on clay soils; tolerate moderate alkalinity. Salt tolerance is higher during germination than during subsequent development. At 20.0 mmhos/cm salinity, seed yields are reduced 50%. Greenhouse studies suggest that tolerance to 3.9 mmhos is more realistic. Lentils are quantitative long-day plants, some cultivars tending to be day-neutral. Lentil is reported to require environments ranging from cool temperate steppe to wet through subtropical dry to moist forest life zones. It tolerates annual precipitation of 2.8-24.3 dm annual mean temperature of 6.3-27.3°C and pH of 4.5-8.2". (Kay, 1979; Duke, 1981)^[5]

Soil

Lentil prefers a sandy soil in a warm sunny sheltered position, prefers well-drained soil and can grow in nutritionally poor soil. It cannot grow in the shade. It prefers dry or moist soil. Lentils grow on many soil types, ranging from sandy to heavy clay soils, and on a large range of pH (4.5-9), provided that the soils are not saline, waterlogged or subject to flooding (Bejiga, 2006)^[2].

Traditional and Medicinal Use

They are considered to be useful in the treatment of constipation and other intestinal affections. Made into a paste, they are a useful cleansing application in foul and indolent ulcers (Chopra et. al., 1986)^[3].

In India, lentils are poulticed onto the ulcers that follow "smallpox and other slow-healing sores" (Duke, 1981)^[5].

In the 6th century, chickpeas were believed to be an aphrodisiac; while curiously enough, lentils were considered to have the opposite effect, and this was probably the reason why the lentil was included in the diet in monasteries on meatless days (Van der Maesen, 1972)^[12].

The primary product is the seed which has a relatively higher contents of protein, carbohydrate and calories compared to other legumes and is the most desired crop because of its high average protein content and fast cooking characteristic in many lentil producing regions (Muehlbauer et al., 1985).

Husks, dried leaves, stems, fruit walls and bran (residues), can be fed to livestock. Lentil residues contain about 10.2% moisture, 1.8 % fat, 4.4% protein, 50% carbohydrate, 21.4% fiber, and 12.2% ash (Muehlbauer et al., 1985).

Chemical Composition

Protein concentration of lentils reportedly range from 22-34.6%, and 100 g of dried seeds contain 340-346 g calories, 12% moisture, 20.2 g protein, 0.6 g fat, 65.0 g total carbohydrate, about 4 g fiber, 2.1 g ash, 68 mg Ca, 325 mg P, 7.0 mg Fe, 29 mg Na, 780 mg K, 0.46 mg thiamine, 0.33 mg riboflavin, 1.3 mg niacin (Adsule et al., 1989; Muehlbauer et al., 1985). Among the cool season legume crops, lentil is the richest in the important amino acids (lysine, arginine, leucine, and sulphur containing amino acids) (Williams et al., 1994). The starch content ranges from 35-53% in the seed and 42% in dry matter while amylose varies from 20.7 to 38.5% of the seed starch (Huisman and Van der poel, 1994; Hulse, 1994)^[7, 8]. "One hundred grams of decorticated lentil seed contain 344 calories, 9.9 % moisture, 25.8 g protein, 1.8 g fat, 58.8 g total carbohydrate, 0.9 g fiber, 3.7 g ash, 24 mg Ca, 271 mg P, 10.6 mg Fe, 0.47 mg thiamine, 0.21 mg

riboflavin, and 1.5 mg niacin. Lentils are a good source of B vitamins, containing per 100 g: 0.26 mg thiamine, 0.21 mg riboflavin, 1.7 mg nicotinic acid, 223 mg choline, 107 mg folic acid, 130 mg inositol, 1.6 mg pantothenic acid, 13.2 mg biotin, and 0.49 mg pyridoxine. Vitamins, except folic and pantothenic acids, increase markedly during sprouting. Dry lentil husks contain 11.1% protein (1.3% digestible), 0.7% fat, 47.5% carbohydrate, 25.6% fiber, and 3.1% ash" (Duke, 1981)^[5]. "About 90% of lentil protein is found in the cotyledons with albumins and globulins being the major fractions. Digestibility coefficients for lentil are relatively high and range from 78-93%, while biological values range from 32-58%. Oleic, palmitic and linoleic are the dominant fatty acids" (Hulse, 1994)^[8].



Fig 1: Lentil seeds



Fig 2: Lentil crop

References

1. Adsule RN, Kadam SS, Leung HK. Lentil. In: CRC hand book of world food legumes (eds. D.K. Salunkehe and S.S. Kadam). Boca Raton, Florida, USA: CRC Press, 1989.
2. Bejiga G. *Lens culinaris* Medik. Record from Protabase. Brink, M. & Belay, G. (Editors). PROTA (Plant Resources of Tropical Africa / Resources végétales de l'Afrique tropicale), Wageningen, Netherlands, 2006.
3. Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants (Including the Supplement). Council of Scientific and Industrial Research, New Delhi, 1986.
4. Cubero JI. "Origin, taxonomy and domestication," in *Lentils*, eds C. Webb and G. Hawtin (London: Commonwealth Agricultural Bureaux), 1981, 15-38.

5. Duke JA. Handbook of legumes of world economic importance. Plenum Press, New York, 1981, p. 52-57.
6. Facciola S. Cornucopia: a source book of edible plants. Published by Kampong Pubns, Vista, CA, 1990.
7. Huisman J, van der Poel AFB. Aspects of the nutritional quality and use of cool season food legumes in animal feed. pp. 53-76. In: F.J. Muehlbauer and W.J. Kaiser (eds.), Expanding the Production and Use of Cool Season Food Legumes. Kluwer Academic Publishers. Dordrecht, The Netherlands, 1994.
8. Hulse JH. Nature, composition and utilization of grain legumes. p.11-27. In: ICRISAT. Uses of tropical grain legumes: Proceedings of a consultants' meeting, 27-30 Mar, 1989. ICRISAT Center, India, Patancheru, A.P. 502 324. India: ICRISAT, 1990.
9. Kays D. Food Legumes. Tropical development and research institute (TPI). TPI Crop and Product Digest, 1979; (3):48-71.
10. Lev-Yadun S, Gopher A, Abbo S. The cradle of agriculture. Science, 2000; 288:1602-1603.
11. Muehlbauer FJ, Cubero JI, Summerfield RJ. Lentil (*Lens culinaris* Medic.). p. 266-311. In: R.J. Summerfield and E.H. Roberts (eds.), Grain Legume Crops. Collins, 8 Grafton Street, London, UK, 1985.
12. Van der Maesen LJG. Cicer L., A monograph of the Genus, with special reference to the chickpea (*Cicer arietinum* L.), its ecology and cultivation, Commun. Agric. University. Wageningen, 1972.
13. Williams PC, Bhatti RS, Deshpande SS, Hussein LA, Savage GP. Improving nutritional quality of cool season food legumes. p. 113-129. In: F.J. Muehlbauer and W.J. Kaiser (eds.), Expanding the Production and Use of Cool Season Food Legumes. Kluwer Academic Publishers, Dordrecht, the Netherlands, 1994.
14. Xu B, Chang SK. Phenolic substance characterization and chemical and cell-based anti-oxidant activities of 11 lentils grown in the Northern United States. Journal of Agricultural and Food Chemistry, 2010; 58:1509-1517
15. Zohary D. Monophyletic vs. polyphyletic origin of the crops on which agriculture was founded in the Near East. Genet. Resour. Crop Evol, 1999; 46:133-142.
16. Nasr ME, Zinhoum RA, Lotfy K. Efficacy of cold plasma against three of stored grain insects. Int J Entomol Res. 2020;5:113-7.