

ISSN Print: 2664-844X
 ISSN Online: 2664-8458
 Impact Factor: RJIF 5.6
 IJAIFS 2023; 5(1): 01-07
www.agriculturaljournals.com
 Received: 05-10-2022
 Accepted: 06-11-2022

Hassan I Mohammed
 Department of Agricultural
 Engineering, College of
 Agricultural Studies, Sudan
 University of Science and
 Technology, Khartoum, Sudan

AbdI Karim D Elffadil
 Department of Agricultural
 Engineering, Faculty of
 Agricultural Sciences,
 University of Gezira, Wad
 Madani, Sudan

Ali Hassan
 Department of Agricultural
 Engineering, Faculty of
 Agricultural Sciences,
 University of Gezira, Wad
 Madani, Sudan

Omer A Abd Alla
 Department of Agricultural
 Engineering, Faculty of
 Agriculture, University of
 Khartoum, Sudan

Haithum R Ramlawi
 Center of Dry land Farming
 Research and Studies, Faculty
 of Agricultural and
 Environmental Sciences, Univ.
 of Gadaref, Sudan

Corresponding Author:
Hassan I Mohammed
 Department of Agricultural
 Engineering, College of
 Agricultural Studies, Sudan
 University of Science and
 Technology, Khartoum, Sudan

Effects of micro-water harvesting techniques on sesame growth, yield and soil moisture on clay soil in rain fed area

Hassan I Mohammed, AbdI Karim D Elffadil, Ali Hassan, Omer A Abd Alla and Haithum R Ramlawi

DOI: <https://doi.org/10.33545/2664844X.2023.v5.i1a.107>

Abstract

Sesame (*Sesamum indicum* L.) is the major cash crop yearly grown in rain fed areas as alternative to Sorghum crop, but their grain yield is remarkably reduced by soil moisture stress when cultivated in cracking clay soil under semi-arid climate. This require employing good Vertisol water management practices that conserve moisture during dry spell period and excess moisture during wet periods. Possible practice to adopt is to utilize relevant water harvesting technique. Therefore, the main objective of this study is to investigate the performance of three water harvesting methods namely: semi-circular bund, ridge and furrow, and sowing on flat soil surface, on soil moisture distribution and Sesame crop growth attributes (Days to 50% flowering, days to 90% maturity, number of branches per plant, and plant height). Yield Attributes (number of seeds per capsule, thousand seed weight,) and seed yield (Weight per unit area). The field study was conducted at El Migrih in Butana rain fed area during 2019/2020 and 2020/2021 growing seasons in vertisol soil type. The experiment was laid out in randomized complete block design (RCBD) with three replications using variety Khidir. Statistical analysis revealed that water harvesting techniques had a significant effect on soil moisture and agronomic parameters. The ridge and furrow method conserved best soil moisture resulting in the highest sesame seed yield (548 kg ha⁻¹) while the lowest yield (390 kg ha⁻¹) was obtained with flat traditional method.

Keywords: Water harvesting methods, soil moisture conservation, vertisol, sesame, rain fed

1. Introduction

Sesame is produced as main cash crop mainly in the rain-fed areas within the central clay plains of Sudan where research is limited by funding and staff. This producing area (2.4 million hectares annually in Northern Gadaref "300-400 mm rain", Central Gadaref "400-500 mm rain") has limited rainfall which is erratic in nature and of uneven distribution which results in soil moisture stress when accompanied by high evaporation. Annual yield of sesame (0.274t/ha) are by far below the international average (0.512 t/ha). Available data showed that yields of sesame progressively declined with time (Elleuch *et al.*, 2011) [7]. Similarly, moisture stress could have resulted when transpiration of water exceeding the rate of absorption. Sesame is planted, annually, in about (2.4 million hectares). The Central Clay Plain is part of the guild of east Africa and dominated by vertisols are renowned by their high difficulties in water management for the soil is hard when dry and sticky when it is wet. Sesame grain yield is easily affected by soil moisture stress in semi-arid climate. Sarhadi, and Sharif, (2014) [22] and Nimir, (2002) [24] both confirm Sesame sensitivity to water shortage by studying effect of deficit irrigation on sesame growth, yield and yield components in drought conditions on base of sustainable agriculture. Goitom *et al.* (2021) [9] studied the effect of land configuration ways (flat, ridge and furrow, and bed furrow) on soil moisture and sesame seed yield. Girma. (2019) [10]. conducted a factorial experiment to study the performance of sesame under different supplementary irrigation applications and nitrogen fertilizer rates. Data on plant height, number of capsules per plant, number of seeds per capsule, thousand seed weight, harvest index, aboveground biomass and grain yield were recorded and used as evaluation indicators (Loggale. 2018) [14]. Sesame crop is characterized by its high sensitivity to water stress at its early growth phases (germination and seedling - Boureima *et al.*, 2011) [2] and at reproductive stage (Nilanthi *et al.* 2015) [18].

Consequently, searching for relevant water harvesting techniques and cultural operations that conserve moisture during dry period and drain excess moisture during wet periods is critically needed. Therefore, in this study, three land water harvesting techniques (WHTs) are proposed as ways for land configuration and to be compared on their effect on sesame productivity. These are: semi-circular bund, ridge and furrow, and sowing on flat. Although other cultural practices such as weeding fertilization and tillage have a great influence on sesame crop performance and yield they will be kept consistent and same in all conducted trials.

2. Materials and Methods

2.1. Study Area

The experiment was conducted in 2019/2020 and 2020/2021 growing seasons at El Gadambalyia in Butana rain fed area located between (latitudes 13°45' and 14 ° 15' N and longitudes 34° 45' and 35°) The soil is cracking vertisol which is sticky when wet and hard when dry with low infiltration rate (2-3 mm/hr), low nitrogen content (0.212%), low organic matter (1.4%) (El Abbas, *et al.*, 2017). The ranges of field capacity, wilting point, and bulk density were 31.69–32.38%, 21.20–21.53%, and 1.31–1.37 gm / cm respectively, and water holding capacity is high (Table 1.0).

Table 1: Physical properties of the soil of the study area

Layer thickness (cm)	Field capacity (%)	Wilting point (%)	Bulk density (gm/cm ³)	EC (dS / m)	pH (%)	Lime (%)	Clay (%)	Sand (%)	Silt (%)	Texture
0 - 30	31.94	21.53	1.31	0.200	8.7	20.22	61.64	8.85	29.51	Clay
30 - 60	31.69	21.20	1.32	0.165	7.7	20.03	66.06	8.56	25.35	Clay
60 - 90	32.38	21.53	1.37	0.156	7.5	19.45	66.98	8.76	24.26	Clay

The annual rainfall ranges between 400 to 500 mm and most of the rainfalls from June to September. The annual mean temperature is (40 °C in summer to 16 °C in winter with

erratic rainfall (350 to 550 mm/annum) (Table 2). The study area is a vast plain of low land suitable for large scale semi-mechanized crop production.

Table 2: The trial was located in an arid zone; where winter is mild and the summer is dry

Climate Element	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Min Temp (C)	16.81	18.87	21.80	24.52	26.37	27.11	24.79	23.92	24.63	24.00	21.05	17.68
Average Temp (C)	24.54	27.00	30.18	33.22	34.18	33.43	29.82	29.23	31.89	31.94	28.75	25.20
Max Temp (C)	32.26	35.12	38.56	41.91	41.99	39.74	34.85	34.54	39.14	39.87	36.44	32.72
RH (%)	14.89	10.37	7.42	6.68	11.53	17.25	25.96	28.73	21.86	14.73	12.70	15.96
Wind (m/s)	1.94	2.04	2.03	1.77	1.64	2.19	2.28	2.04	1.66	1.42	1.83	1.87
RF (mm)	0.02	0.00	0.60	1.49	12.38	18.75	76.75	100.48	28.83	9.15	2.54	0.00
Solar	20.69	23.61	26.17	27.42	25.09	24.16	20.30	20.07	21.33	20.94	20.80	19.46

2.2. Experimental treatments, design and cultural practices

The experiment was made in a randomized complete block design (RCBD) with three replications. Treatments includes: semi-circular bund, ridge and furrow, and sowing on flat soil surface. The dimensions of the semi-circular bund (Critchley and Siegert, 1991) ^[4] are: height: 0.2 to 0.5 m, base width : 0.70 m, side slopes : 3 : 1, crest width: 0.1 - 0.25 m, and diameter: 12 m - 40 m. The ridge and furrow are with a furrow depth of 30 cm; furrow spacing at 0.8 m and bed height is 30 cm from soil surface. The flat land sowing is typical to farmer practice using wide level disc machine. These water harvesting techniques were intended to configure the land for maximizing rainfall infiltration, minimizing erosion, and reduction of total runoff to facilitates drainage and finally improves water use efficiency. A spacing distance of 1.5m was kept between blocks between and 1.0 m between plots inside each block. Khidir sesame variety was sown after field was plowed with disc harrow, leveled and shaped according to specifications of the intended water harvesting technique. Other cultural practices (seed rate, weeding, fertilizer) as recommended by Agricultural Research Corporation (ARC) is kept the same for all treatments (Ahmed, 1998, Loggale. 2018) ^[1, 14].

2.3. Data Collection

Data collected includes crop data (growth stages, Yield and yield components), and Soil moisture data).

2.3.1. Crop data

Crop growth stages were determined by visual observation and guided by Doorenbos *et al.* (1979) ^[3].

1. Initial stage (from planting to initiation of seedlings and ends after 15 days from sowing);
2. Establishment of Seedlings to vegetative stage (after 30 days from sowing)
3. Vegetative stage (after 50 days from sowing);
4. Flowering stage (after 70 days from sowing and when 50% of plants flower);
5. Ripening or Maturity stage (after 90 days when (90%) of the crop leaves turned yellow and most of the lower capsules started to open).

The crop growth components (number of branches per plant, number of capsules per plant, number of seeds per capsule and plant height) were determined from randomly selected five plants per plot. Crop growth rate was determined by counting days to 50% flowering and days to 90% maturity. The yield components (number of seeds per capsule, thousand seed weight, and seed yield as weight per area) were determined after harvesting: For number of seeds per capsule, the seeds of three capsules (lower, medium, and uppermost position on the plant) from each of five plants were counted. 1000 seeds number and weight were determined from each plot after sun drying. Seed yield per plot was weighed in grams per plot unit area (kg/ha).

2.3.2. Soil moisture data

The gravimetric oven dry method was used to determine soil moisture from a profile of one meter depth at interval of 0.2m using an auger. Data was replicated three times per plot, at two weeks interval. Soil data were taken after 15, 30, 45, 60, and 75 day from sowing.

2.4. Data Analysis

Collected data was analyzed by descriptive statistics using Statistic 10 software to determine analysis of variance using the least significant difference of 5% and, mean separation using least significance difference (DB Duncan, 1955) [5].

3. Results and Discussion

3.1. Impacts of water harvesting techniques on crop growth and yield

3.1.1 Crop Growth Attributes

Days to flowering, and to maturity, number of branches per plant, and plant height (cm) crop growth attributes are given in table 1.0. The table shows their results for analysis of variance, coefficient of variation (Cv %) and, mean separation using LSD. (Girma. 2019, and Goitom *et al.* 2021) [10, 9]

Table 1: Crop Growth Attributes (Days to flowering, and maturity, number of branches per plant, and plant height) for the different WHTs

Treatment	Days to 50% flowering	Days to 90% Maturity	No. Braches/plant	Plant Height(cm)
Semi-circular	42	93 a	2.7	78 a
Flat	43	94 b	2.6	69b
Ridge and Furrow	41	92 a	3.0	75a
Cv. %	2.3	1.0	7.5	6.1
LSD. (0.05)	3	1.9	0.7	0.9

a- Days to Flowering: Table 1.0 shows analysis of variance for days to flowering determined for the various WHTs and illustrated an insignificant effect ($p>0.05$) of WHT on the 50% flowering. The ridge and furrow and took shorter period (41 days) for the 50% flowering, while the longer days to the 50% flowering

were needed for flat land sowing. Conservation and availability of moisture retained by ridge and furrows may be the cause of speeding flowering processes leading to early maturing (Figure 1). This result agrees with Reddy *et al.*, (2009) [20] and Goitom *et al.* (2021) [9].




Fig 1: Days to 50% flowering achieved by various WHTs

b- Days to 90 % Maturity: Figure 2.0 illustrate the results of days to 90 % Maturity reached by the studied WHTs. Early maturity as proxy-indicator of healthy growth is achieved by ridge and furrow, followed by semi-circular bund, and the latest is sowing on flat soil surface. This result is in line with days to flowering (Figure1.0).The differences between ridge and furrow, and semi-circular bund are not significant ($p<0.05$) but they both differ significantly from sowing on flat (Table 1.0).

c- Number of branches per plant: The data in table1.0 indicated that the all WHTs didn't show a significant difference ($p>0.05$) on the number of branches per plant. The highest number of branches (3.0) was recorded by ridge and furrow and the lowest (2.6) was recorded under the flat land WHT (Figure 3.0). This is may be taken as proxy – indicator of more yield (Goitom *et al.* 2021) [9]




Fig 2: Days to 90 % Maturity reached by the studied WHTs




Fig 3: Number of branches per plant developed by the studied WHTs.

d- Plant height (cm): As given in Table 1.0 it is evident that plant height showed significant differences at ($p < 0.05$) between WHTs treatments. The minimum plant height (69 cm) was recorded on the flat land while that measured by Semicircular and Plant Height are higher but not significantly ($p > 0.05$) different from each other (Figure 4). This result is typical to the results obtained for days to maturity. Plant height is a yield component that is affected by, pests and disease, and weed infestation, and stress by in excess or deficit water. It is evident from figure 4 that capability of WHT to conserve soil moisture can be expressed by plant height and is rank the WHTs in a descending order as Flat (69 cm), Ridge and Furrow (75 cm), and Semicircular (78 cm). This result is the same to that stated by Goitom *et al.* (2021) [9].




Fig 4: Recorded plant height (cm) with the three WHTs

3.1.2 Yield and Yield Attributes:

In this study number of capsules per plant, their respective number of seed per capsule and weight of seed, are taken as essential factors in determining the final yield obtained by the crop. The number of these capsules is function of rise or drop in moisture status. Short of water affect translocation of assimilates and nutrients and flooding reduce initiation, setting and development of flowers. (Kadkhodaie *et al.*, 2014; Hailu *et al.*, 2018, Goitom *et al.* 2021) [12, 11, 9]. Consequently, the performance of the three WHTs with respect to yield and yield attributes (No Capsule per plant, No Seeds per Capsule, Seed weight, and Grain Yield) is depicted in table 2.0.

Table 2: Yield and Yield Attributes obtained by WHTs

Treatment	No Capsule / plant	No Seeds / Capsule	Seed weight	Grain Yield Kg / ha
Semicircular	32.5 ab	41 a	3.3 a	454 b
Flat	29.3 b	39 a	3.2 a	390 c
Ridge and Furrow	33.9 a	43 a	3.4 a	548 a
Cv%	7.5	4.9	3.0	17.0
LSd (0.05)	9.5	4.5	0.3	23.7

i- Number of capsules per plant: Table 2.0 showed significant differences ($p < 0.05$) on capsules number per plant with different WHTs. The highest number of capsules (33.9) was recorded at the ridge and furrow WHT and it is significantly higher than measured in flat sowing but do not differ significantly from capsules of semi-circular bund WHT. However, Malik *et al.*, (2003) [17], and Goitom *et al.* (2021) [9] gave similar results.




Fig 5: Number of capsule per plant for the studied WHTs

ii- Number of seeds per capsule: Table 2.0 shows that there is no significant difference ($p > 0.05$) in number of seeds per capsule recorded for the three WHTs. Figure 6 shows that the highest number of seeds per capsule is found with Ridge and furrow method, followed by semi – circular bund and this result is in agreement with Malik *et al.*, (2003) [17] and Goitom *et al.* (2021) [9].




Fig 6: Number of seeds per capsule for the studied WHTs

iii- Thousand Seed Weight: The analysis of variance showed insignificant difference ($p > 0.05$) between WHTs for thousand seed weight (Table 2). According to figure 7 the highest thousand seed weight (3.4 g) was recorded at the ridge and furrow, followed by semi – circular bund (3.3 g) and the least is noted for sowing on flat (3.2 g). This is in line with Malik *et al.*, (2003) [17] and Goitom *et al.* (2021) [9].




Fig 7: Thousand seed weight obtained by WHTs

iv- Final grain yield (Kg/ha): Impact of WHT on grain yield given in table 2.0 indicates that there is significant difference ($p < 0.05$) in the yield obtained by WHTs. The highest yield (548 Kg/ha) was measured for Ridge and Furrow followed by semi – circular bund (548 Kg/ha) and then sowing on flat (390 Kg/ha). In a similar study Goitom *et al.* (2021) ^[9] reported that the highest sesame seed yield (558 kg ha) was recorded at ridge and furrow, while the lowest (400 kg ha⁻¹) was found at the flat land configuration method.

The high yield obtained by Ridge and furrow method and the minimum yield with conventional application method is in agreement with Hailu *et al.*, (2018) ^[11] and Goitom *et al.* (2021) ^[9]. The yield values for sowing on flat is above the average yield recorded for the central clay plains of Sudan (274 Kg/ha), while the yields recorded for Ridge and Furrow and semi – circular bund are higher than the international average (512 Kg/ha). Reddy *et al.*, (2009) ^[20], Goitom *et al.* (2021) ^[9] and Rajput *et al.*, (2009) ^[19] attributed the highest yield values obtained under the Ridge and Furrow to be due to the high efficiency build in the water harvesting techniques to conserve soil moisture resulting in higher plants height and more branches, large number of capsules per plant, and number of seeds per capsule.




Fig 8: Final grain yield (Kg/ha) obtained by WHTs

3.2. Impacts of WHTs on soil moisture at crop growth stages

Table 3 shows soil moisture retained by WHTs during different growth stages. The table indicate that there is no significant differences ($p < 0.05$) in measured soil moisture at 15 days from sowing (during initial stage) and at 50 days

from sowing (vegetative stage) between the different treatments of WHTs. In contrast, the table shows significant difference ($p < 0.05$) in the measured soil moisture during seedling (50 DAS), flowering (70 DAS), and ripening (90 DAS) stages. The amount of soil water stored by each WHT is attributed to difference in their capability to retain soil moisture (Tantawy *et al.*, 2007 and Goitom *et al.* 2021) ^[23, 9]. During the growth period of vegetative growth to flowering (50 DAS) and in the growth period of flowering to grain filling stage (70 DAS) the WHTs there significant differences in WHTs of store water with least amount is retained by fat soil surface (its shape provide no storage facility). During grain filling stage (90 DAS) the high water stored by both ridge and furrow and semi-circular bund resulted in generating high crop grain yield. This result agrees with Reddy *et al.*, (2009) ^[20] who claimed that the maximum yield obtained by the ridge and furrow technique is attributed to the technique capability to store high water reserve. Likewise in a typical study conducted by Goitom *et al.* (2021) ^[9] indicated that the ridge and furrow methods conserved best soil moisture for highest plant and seed yield performance.

Table 3: Moisture retention by the studied WHTs during crop phenological growth stages

Stage	Initial	Seedlin g	Vegetativ e	Flowerin g	Ripenin g
Date from Sowing	15 DAS	30 DAS	50 DAS	70 DAS	90 DAS
Profile	0 - 90 cm				
WH method	Soil moisture mm				
Semi-circular	48.1	56.8	80.2	65.7	42.9
Flat	47.9	49.7	79.3	64.0	35.4
Ridge & Furrow	50.2	60.5	80.5	68.8	49.9
mean	48.7	55.7	80.0	67	42.7
Cv %	2.7	9.9	0.8	4.0	17
Significance test	Ns	Sd	Ns	Sd	Sd

Ns=not significant; Sd significant

For vertisols of central clay plains of Sudan Farbrother (1987) reported that soil profile moisture capacity is 93 mm, which need to be filled during each irrigation event. As given in Figure 9 shows that the variation of the moisture of the whole soil profile along the length of the growing season follow a curve of bell shape showing lower retention during early stage and increase in water accumulated in the soil profile till the plant reach full canopy and then the stored water starts to diminish by rise of crop water depletion due to increase in crop evapotranspiration (Roa, and Raju, 1991 and Goitom *et al.* 2021) ^[21, 9].




Fig 9: Variation of soil moisture along the length of the growing season by the different WHTs

The insignificant difference ($p>0.05$) on soil moisture content for 15 days after sowing at entire soil depth is related to the rapid water intake into the soil due to effect of cracks irrespective to the shape and configuration of soil surface made by any one the water harvesting technique (Langham *et al.*, 2008, Goitom *et al.* 2021^[15, 9], and Keteku *et al.*, 2020)^[13]. Typical to the results depicted in figure 9 Reddy *et al.*, (2009)^[20] reported the highest moisture conservation efficiency obtained by Ridge and Furrows compared to the semi-circular or flat land configuration at 30, 70 and 90 days after sowing. Likewise, Hailu *et al.*, (2018)^[11] made study of nine treatments with three levels of irrigation water for identifying optimum soil moisture stress for sesame to determine appropriate water-saving irrigation methods and also productivity under limited water resource conditions. Hailu *et al.*, (2018)^[11] concluded that the conventional furrow application method is best practice of soil moisture -saving strategies to improve sesame yield

4. Conclusions

The WHTs have direct and significant impacts on soil moisture and sesame seed yield. The result obtained ranked the studied WHTs on basis of soil moisture retention, yield, yield attributes in descending order of, effectiveness as: ridge and furrow technique, semi-circular bund and sowing on flat soil surface. It is thus recommended to adopt ridge and furrow technique as effective and important climate change adaptation strategy in dry land areas where water shortage, late onset and early cessation of rain occur.

5. Acknowledgment

The financial support provided by Ministry of Higher Education is greatly acknowledged

6. References

- Ahmed MH. Release of cultivar Khidir for production in high rainfall areas, (Southern Gadarif and Damazine) Sudan Journal of Agric. Research. 1998;1(1):89.
- Boureima S, Eyletters M, Diouf M, Diop TA, Van Damme P. Sensitivity of seed germination and seedling radical growth to drought stress in sesame (*Sesamum indicum* L.). Research Journal of Environmental Sciences. 2011;5(6):557-564.
- Doorenbos J, Kassam AH, Bentvelsen CIM, Branscheid V, Plusje JMg A, Smith M, *et al.* Yield response to water. Fao.org. Irrigation and drainage, 1979, 33.
- Critchley Will, and Klaus Siegert. A Manual for the Design and Construction of Water Harvesting Schemes for Plant Production with contributions from: C. Chapman, FAO Project Manager M. Finkel, Agricultural Engineer, Yoqneam, Israel. Fao.org. Food and agriculture Organization of the United Nations - Rome, 1991.
- Duncan DB. Multiple range and multiple F tests. Biometrics. 1955;11:1-41.
- El Abbas Doka MA, Aitken J, Munro NR. Studying the Characteristics of Vertisols to Set Up Field Management Practices at Dinder Area (Sennar State-Sudan). Advances in Image and Video Processing. 2017;5(5):21-21.
- Elleuch M, Bedigian D, Zitoun A. Sesame (*Sesamum indicum* L.) seeds in food, nutrition, and health. In Nuts and Seeds in Health and Disease Prevention. Academic Press, 2011, p. 1029-1036.
- Farbrother HG. Supplementary Irrigation. Management of Vertisols under semi-arid conditions Seminar organized by The International Board for Soil Research and Management (IBSRM). Proceeding No.6. Marc Latham and Peter Ahm and Colin R. Elliott Editors, 2017, P267-285. ISBN 974-7614 – 47 – 2. Printed in Thailand.
- Goitom Teame, Alemtsehay Tsegay, Dawit Weldearegay. Effect of land configuration ways on soil moisture and sesame seed yield in Ethiopia. Teame *et al.*/Turk J. Food Agric. Sci. 2021;3(1):1-6, 2021. - ISSN: 2687-3818. Doi:10.14744/turkjfas.2021.001
- Girma Abadi Berhane. Performance of Sesame (*Sesamum indicum* L.) Under Different Supplementary Irrigation and Nitrogen Fertilizer Levels in Humera, Northern Ethiopia. Mod Concep Dev Agrono, 2019, 5(1). MCDA.000605.2019. DOI: 10.31031/MCDA.2019.05.000605
- Hailu EK, Urga YD, Sori NA, Borona FR, Tufa KN. Sesame Yield Response to Deficit Irrigation and Water Application Techniques in Irrigated Agriculture, Ethiopia. Hindawi International Journal of Agronomy, 2018, Volume 2018, Article ID 5084056, 6 pages <https://doi.org/10.1155/2018/5084056>
- Kadkhodaie A, Morteza Zahedi M, Razmjoo J, Pessaraki M. Changes in some anti-oxidative enzymes and physiological indices among sesame genotypes (*Sesamum indicum* L.) in response to soil water deficits under field conditions. Acta Physiologiae Plantarum. 2014;36(3):641-650.
- Keteku AK, Kadam AK, Dana S, Blege PK. Influence of land configuration and fertilization techniques on soybean (*Glycine max* (L.) Merrill.) productivity, soil moisture and fertility. Acta Agriculture Slovenica. 2020;115(1):79-88.
- Loggale LB. Performance of Two Sesame Cultivars as Influenced by Supplemental Irrigation at Abu Naama." IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS). 2018;11(9):06-11.
- Langham DR, Riney J, Smith G, Wiemers T. Sesame grower guide. Sesaco sesame coordinators, Lubbock, TX Retrieved, 2008. <http://www.sesaco.net>.
- Taminaw Zewdie, Nuru Seid Tehulie. Review on effects of plant densities and nitrogen fertilization on sesame (*Sesamum indicum* L.) Yield and yield components. Int. J Res Agron 2019;2(1):42-47. DOI: 10.33545/2618060X.2019.v2.i1a.66
- Malik MA, Saleem MF, Cheema MA, Ahmed S. Influence of different nitrogen levels on productivity of sesame (*Sesamum indicum* L.) under varying planting patterns. International Journal of Agriculture Biology. 2003;5(4):490-492.
- Nilanthi D, Alawathugoda C, Ranawake A. Effects of water stress on yield and some yield components of three selected oil crops; Groundnut (*Arachis hypogea* L.), sunflower (*Helianthus annuus* L.) and sesame (*Sesamum indicum* L.) International Journal of Scientific and Research Publications. 2015;5(2):2250-3153.
- Rajput R, Kauraw DL, Bhatnagar RK, Bhavsar M, Velayutham M, Lal R. Sustainable management of vertisols in central India. Journal of Crop Improvement. 2009;23(2)119-135.

20. Reddy BS, Maruthi V, Adaje RV, Madal UK. Effect of different land configuration practices on productivity of sorghum–pigeon pea intercropping system in Shallow, 2009.
21. Roa V, Raju C. Effects of soil moisture stress at deferent development phases on growth and yield of sesame, Journal of Oilseeds Research. 1991;8(2):240–243.
22. Sarhadi J, Sharif M. effect of deficit irrigation on sesame growth, yield and yield components in drought conditions on base of sustainable agriculture, International Journal of Farming and Allied Sciences. 2014;3(10):1061–1064.
23. Tantawy MM, Oudu SA, Khalil FA. Irrigation optimization for different Sesame varieties grown under water stress condition, Journal of Applied Science Research. 2007;3(1):7-12.
24. Nimir ESN. Effect of water stress applied at different stages of growth on the performance of two sesame (*Sesamum indicum* L.) cultivars. M.Sc. Thesis. Fac. of Agric. University of Khartoum; c2002.