

ISSN Print: 2664-844X
ISSN Online: 2664-8458
NAAS Rating (2025): 4.97
IJAFA 2025; 7(8): 1342-1346
www.agriculturaljournals.com
Received: 21-05-2025
Accepted: 24-06-2025

PH Kare

Department of AHDS, Post
Graduate Institute MPKV,
Rahuri, Maharashtra, India

BD Patil

Department of AHDS, Post
Graduate Institute MPKV,
Rahuri, Maharashtra, India

DK Kambale

Department of AHDS, Post
Graduate Institute MPKV,
Rahuri, Maharashtra, India

SS Dighe

Department of Horticulture,
Post Graduate Institute
MPKV, Rahuri, Maharashtra,
India

SD Shinde

Department of Statistic, Post
Graduate Institute MPKV,
Rahuri, Maharashtra, India

Corresponding Author:

PH Kare

Department of AHDS, Post
Graduate Institute MPKV,
Rahuri, Maharashtra, India

Process optimization for preparation of *lassi* blended with grape juice (*Vitis vinifera*)

PH Kare, BD Patil, DK Kambale, SS Dighe and SD Shinde

DOI: <https://www.doi.org/10.33545/2664844X.2025.v7.i8m.710>

Abstract

Lassi was prepared by using cow milk with optimized levels of grape juice and sugar. The sensory evaluation of *lassi* was done by semi trained judges. The investigation was undertaken to explore the possibilities of utilizing grape juice in *lassi* manufacture to improve the health benefits of product. The product obtained was subjected for chemical analysis. The *lassi* sample (T₄) prepared with 6 per cent grape juice and 15 per cent sugar level received the highest overall acceptability score i.e., 8.47. The sensorily most acceptable treatment combination (T₄) has an average chemical composition of 3.09% fat, 3.26% protein, 78.08% moisture, 21.92% total solids, 0.75% ash, 0.85% acidity, 4.15 pH, 4.34% reducing sugars, 10.48% non-reducing sugar and 14.82% total sugars.

Keywords: Grape juice, cow milk, sugar, *Lassi*

Introduction

Fermented milk and its products are vital components of human nutrition, offering a range of health benefits. These products retain all the essential nutrients of milk while improving their bioavailability. During fermentation, milk proteins coagulate, making them easier to absorb and digest particularly beneficial for vulnerable groups such as children, the elderly, and individuals with gastric ulcers (Singh and Amin, 1983) ^[19]. *Lassi* is a traditional fermented dairy drink commonly enjoyed in many cultures. It is prepared by stirring whole curd into a delicious drink with addition of sugar or salt a small amount of cold water or ice to make the product flowable (Kedaree *et al.*, 2021) ^[11]. This beverage has historically been an important part of diets in different regions, valued both for its refreshing qualities and nutritional benefits.

Grape juice contains high amounts of natural sugars, vitamins, and polyphenols so it can be used as a natural sweetener and functional ingredient. Grape is one of the largest fruit crops grown worldwide and rich in carbohydrates, organic acids, amino acids, vitamins, melatonin and phenolic compounds (Zare and Lashkari, 2021; Kersh *et al.*, 2023) ^[23, 12]. The grape berry and its derived products have potent antioxidant, anticarcinogenic, antibacterial, antidiabetic, and anti-inflammatory activities as well as cardioprotective, hepatoprotective and neuroprotective effects many people are becoming aware of the importance of their consumption in their daily life due to positive health benefits (Bendaali *et al.*, 2022) ^[3]. Grapes are rich in bioactive compounds such as phenolic acids, flavonoids, anthocyanin, stilbenes, and lipids (Shiraishi *et al.*, 2010) ^[20].

Materials and Methods

The following materials were used for the present investigation.

Collection of Cow Milk

Fresh clean, composite sample of crossbreed cow milk were obtained from Research Cum Development Project on Cattle, Department of Animal Husbandry and Dairy Science, MPKV, Rahuri.

Collection of Grape cultivar

Good quality fresh grapes fruits of cultivar Concord, Arka Shyam and Bangalore Blue were procured from AICRP on fruits MPKV, Rahuri and grape juice extraction was done.

Microbial Cultures

The standard dahi culture i.e., LF 40 containing *Lactobacillus* spp. were used in this study and was procured from NDRI, Karnal.

Chemicals

Analytical (AR) or guaranteed grade (GR) reagents were used in the chemical analysis.

Methods Phase –I Preliminary Trials

Preliminary trials were conducted to optimize grape cultivar, juice levels and sugar levels.

Phase –II Experimental Trails

On the basis of results of sensory evaluation of preliminary trials 3, 6 and 9% juice of concord cultivar and 12 and 15% sugar were used to prepare the lassi samples.

Preparation of lassi sample

Lassi was prepared as per the method suggested by De, (2004) with slight modifications.




Fig 1: Flow chart for preparation of lassi

Procedure: *Lassi* was prepared as per the procedure described by De, (2004) [4].

Treatment combinations

For preparation of *lassi* by using grape juice (*Vitis vinifera*), the treatment combinations were finalized in preliminary trials as per follows:

Treatment's No.	Treatment combination
T ₀	Dahi + 12% sugar
T ₁ (J1S1)	Dahi + 3% grape juice + 12% sugar
T ₂ (J1S2)	Dahi + 3% grape juice + 15% % sugar
T ₃ (J2S1)	Dahi + 6% grape juice + 12% sugar
T ₄ (J2S2)	Dahi + 6% grape juice + 15% sugar
T ₅ (J3S1)	Dahi + 9% grape juice + 12% sugar
T ₆ (J3S2)	Dahi +9% grape juice + 15% sugar

Table 1: Sensory score of fresh *lassi* samples (Sensory score out of 9)

Treatments	Flavour	Colour and Appearance	Consistency	Overall Acceptability
T ₀	7.83 ^e	7.80 ^f	7.90 ^e	7.84 ^f
T ₁	8.01 ^d	8.02 ^d	8.10 ^d	8.04 ^d
T ₂	8.20 ^c	8.17 ^c	8.25 ^c	8.20 ^c
T ₃	8.33 ^b	8.34 ^b	8.30 ^b	8.32 ^b
T ₄	8.46 ^a	8.56 ^a	8.40 ^a	8.47 ^a
T ₅	7.94 ^d	7.92 ^e	7.85 ^f	7.90 ^e
T ₆	7.75 ^f	7.88 ^e	7.76 ^g	7.79 ^g
S.E.(m) ±	0.02	0.03	0.004	0.003
CD at (5%)	0.07	0.09	0.012	0.011

Flavour

The mean flavour score of *lassi* for treatments T₀, T₁, T₂, T₃, T₄, T₅, and T₆, were 7.83, 8.01, 8.20, 8.33, 8.46, 8.10 and 7.95 respectively. Flavour scores were significantly ($p < 0.05$) affected by both ingredients. Treatments T₁ and T₅ were at par with each other. Treatment T₄ achieved the highest flavour score of 8.46.

The flavour results from this study were similar to those reported by Gaikwad *et al.* (2018) [6], they prepared a menthol juice *lassi* and found an average sensory score for flavour ranging from 7.31 to 8.80. Roy *et al.* (2019) [18] studied that flavour of the four samples of grape juice yoghurt got near about same score ranging from sample S1 7.17, S2 7.33, S3 7.17 and S4 7.17.

Colour and appearance: For treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆, the *lassi* received colour and appearance scores of

7.80, 8.02, 8.17, 8.34, 8.56, 7.92 and 7.88 respectively. Colour and appearance scores were significantly ($P < 0.05$) affected by both ingredients. Out of all the treatments, treatment T₄ received the highest score (8.56), while T₀ shows lowest score (7.80). Treatments T₁, T₂, T₃, T₄ and T₅ shown notable variations from one another.

The findings of this study align closely with those reported by Avtade (2010) [1] and Ghule *et al.* (2015) [7], they noted that the scores for the colour and appearance of *lassi* improve with increasing levels of pomegranate juice and strawberry pulp, up to a certain point, after which they gradually decline.

Consistency

The consistency score for treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were as 7.90, 8.10, 8.25, 8.30, 8.40, 7.85 and 7.76 respectively. The highest consistency score was observed for treatment T₄ (8.40), while the lowest score was noted for treatment T₆ (7.76).

The results obtained in this study are in agreement with the results obtained by Dhumal *et al.* (2018) [5] who reported that *lassi* prepared from 15 per cent muskmelon and 2.5 per cent pudina extract was superior in consistency than other treatments.

Overall acceptability

The scores of the *lassi* samples in terms of overall acceptability for treatment T₀, T₁, T₂, T₃, T₄, T₅, and T₆ were 7.84, 8.04, 8.20, 8.32, 8.47, 7.90 and 7.79 respectively. The formulation including 6 per cent grape juice and 15 per cent sugar received the highest score, whereas the control sample, which contained no grape juice, received the lowest score.

Compared to all other treatments, treatment T₄ demonstrated the best colour, flavor, consistency and was considerably ($p < 0.05$) superior.

Navale (2020) [15] observed that increase in the level of orange juice resulted in better overall acceptability score of *lassi* up to a certain limit and there after it decreases proportionately.

Chemical composition of fresh *lassi* samples

The freshly prepared *lassi* samples were subjected to chemical analysis *viz.* fat, protein, moisture, ash, acidity, reducing sugars, non-reducing sugars, total sugars, total solids and pH value. The observations and corresponding statistical analysis are presented in table 2.

Table 2: Chemical composition of fresh *lassi* samples

Treatment Combination	Fat (%)	Protein (%)	Total solids (%)	Moisture (%)	Ash (%)	Acidity (%)	pH (%)	Reducing Sugar (%)	Non- reducing Sugar (%)	Total Sugar (%)
T ₀	3.30 ^a	3.34 ^a	21.09 ^g	78.91 ^a	0.60 ^f	0.72 ^f	4.31 ^a	3.65 ^d	10.20 ^g	13.85 ^g
T ₁	3.25 ^b	3.32 ^b	21.47 ^f	78.53 ^b	0.63 ^e	0.76 ^e	4.27 ^b	4.01 ^c	10.26 ^f	14.27 ^f
T ₂	3.20 ^c	3.30 ^c	21.49 ^e	78.51 ^b	0.70 ^d	0.79 ^d	4.22 ^c	3.99 ^c	10.30 ^e	14.49 ^e
T ₃	3.14 ^d	3.28 ^d	21.90 ^d	78.10 ^c	0.73 ^c	0.83 ^c	4.18 ^d	4.36 ^b	10.39 ^d	14.75 ^d
T ₄	3.09 ^e	3.26 ^e	21.92 ^c	78.08 ^c	0.75 ^c	0.85 ^c	4.15 ^d	4.34 ^b	10.48 ^c	14.82 ^c
T ₅	3.04 ^f	3.24 ^f	22.40 ^b	77.60 ^d	0.79 ^b	0.89 ^b	4.12 ^e	4.73 ^a	10.60 ^b	15.33 ^b
T ₆	2.99 ^g	3.22 ^g	22.48 ^a	77.52 ^e	0.82 ^a	0.92 ^a	4.07 ^f	4.72 ^a	10.73 ^a	15.45 ^a
S.E.(m) ±	0.007	0.003	0.003	0.004	0.004	0.005	0.007	0.005	0.006	0.008
CD @ (5%)	0.021	0.009	0.010	0.013	0.012	0.016	0.021	0.016	0.019	0.024

Fat

The fat content for treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were 3.30, 3.25, 3.20, 3.14, 3.09, 3.04 and 2.99 per cent

respectively. The fat content shows significant difference. This was due to lower per cent of fat in grape juice. The

result of fat content in the present investigation was comparable with the finding following research workers. Jadhav *et al.* (2014) ^[14] observed that the fat percentage of the lassi was decreased from 3.25 per cent to 2.92 per cent from treatment T₀ to treatment T₄. The level of fat per cent was decreased as per increase in level of citrus juice in lassi.

Protein

The protein content of lassi significantly ($p < 0.05$) influenced due to addition of grape juice and sugar. The protein content is ranged from 3.34 to 3.22 per cent. The protein content was decreased due to the less amount of protein content in grape juice.

Nur Hossain *et al.* (2012) ^[9], observed that there was a reduction in protein content with the addition of grape juices to yogurt. The above findings are in agreement with the result recorded by Upadhyay *et al.* (2017) ^[22], who reported protein content ranges from 3.53 to 2.82, the protein content in lassi decreased with the increasing level of carrot juice.

Total solids

The total solids content of freshly prepared lassi samples are presented in table 2. The values for the total solids content of lassi were 21.09 to 22.48 per cent for the treatments T₀ to T₆ respectively. The total solids content in the lassi samples varied significantly ($p < 0.05$) with the addition of grape juice.

Tambe *et al.* (2017) ^[21] observed a significant enhancement in lassi's total solids content attributed to the incremental addition of papaya pulp.

Moisture

The data shows that the moisture range of lassi. Treatment T₀, T₁, T₂, T₃, T₄, T₅ and T₆ had moisture content of 78.91, 78.53, 78.51, 78.10, 78.08, 77.60 and 77.52 per cent respectively. Treatment T₀ was the highest moisture content of 78.91 per cent, while T₆ had the lowest i.e., 77.52 per cent. The treatments T₁, T₂ as well as T₃ and T₄ were at par with each other. The moisture content was decreased as grape juice and sugar level increased from treatment T₀ to T₆, this might be due to grape juice had low moisture content as compare to cow milk.

Prabhakar (2018) ^[14] undertook the moisture percentage for various sample of mango fortified lassi which are ranges from 79.87 to 76.05 for treatments T₁ to T₄ and it was observed that moisture percentage decreased from T₁ to T₄.

Ash

The ash content of freshly prepared lassi samples were 0.60, 0.63, 0.70, 0.73, 0.75, 0.79 and 0.82 per cent T₀, T₁, T₂, T₃, T₄, T₅ and T₆ respectively. As the per cent of grape juice increased there is significant ($p < 0.05$) increased in ash content could be due to grape juice contain various nutrients like iron, calcium, zinc, magnesium etc. The treatments T₃ and T₄ were at par with each other.

The results of present study are agreement to the results of Ghule (2015) ^[7] that with increased in the strawberry pulp increased the ash content in lassi.

Acidity (% LA)

The acidity (% LA) of freshly prepared lassi samples were recorded as 0.72% for T₀, 0.76% for T₁, 0.79% for T₂, 0.83% for T₃, 0.85% for T₄, T₅ 0.89% and 0.92% for T₆. The acidity of lassi significantly ($p < 0.05$) influenced due to

addition of grape juice. The acidity in the lassi sample was increased as the level of grape juice increased in the lassi samples because grape juice is acidic in nature. The treatments T₃ and T₄ were at par with each other.

The above results were comparable with the findings of following research workers. Matkar (2010) ^[14] who reported acidity in the range of 0.77 per cent LA (control sample) and 1.04 per cent LA in the lassi blended with papaya pulp. Bagal *et al.* (2018) ^[2] observed that as the amount of papaya pulp increased, there was a corresponding rise in the acidity level of the lassi.

pH

The pH for the treatment T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were 4.31, 4.27, 4.22, 4.18, 4.15, 4.12 and 4.07 respectively. The pH values of lassi samples significantly ($P < 0.05$) declined due to addition of grape juice to lassi. The treatments T₃ and T₄ were at par with each other. The maximum decline in pH value (4.07) was observed in lassi (T₆) prepared using 9 per cent grape juice. The above results were comparable with the findings of following research workers.

These results obtained in present study is in agreement with the results obtained by Ghule (2015) ^[7] strawberry pulp decreases the pH in lassi with increased in strawberry pulp.

Reducing sugars

The reducing sugar content in lassi for treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were 3.65, 4.01, 3.99, 4.36, 4.34, 4.73 and 4.72 per cent, respectively. These variations were statistically significant ($p < 0.05$), demonstrating that the quantity of grape juice significantly influenced the reducing sugar content in the lassi. The treatments T₁, T₂ and T₃, T₄ as well as T₅ and T₆ were at par with each other.

Khore (2013) ^[13] reported that the mean values of reducing sugar in the lassi samples were ranged from 3.82 (T₃) to 3.73 (T₀) percent. It was noticed that as the level of addition of strawberry pulp increased the reducing sugar content in the lassi samples increased significantly. Chakka whey beverage prepared using mango pulp as reported by Hapse (2004) ^[8] had 4.19 percent reducing sugar.

Non-reducing sugar

The Non-reducing sugar content of T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were 10.20, 10.26, 10.30, 10.39, 10.48, 10.60 and 10.73 per cent, respectively. It was observed that the control sample had lowest non-reducing sugar content, T₀ (10.20%), whereas treatment T₆ samples had the highest non-reducing sugar content (10.73%). The data indicated that the non-reducing sugar content in lassi increased with the addition of sugar. Treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were differ from one another considerably ($p < 0.05$).

Similar findings were reported by Hossain (2012) ^[9] the carbohydrate content was highest in G3 (15% grape juice) type yoghurt and the lowest in plain (P) yoghurt. Statistical analysis showed that there were significant differences among the different types of fruit yoghurt in terms of carbohydrate content.

Total sugars

It is revealed that total sugars content in lassi samples of T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were 13.85, 14.27, 14.29, 14.75, 14.82, 15.33 and 15.45 per cent, respectively. Treatment T₀ samples had the lowest total sugars level (13.85%), whereas treatment T₆ samples had the highest content (15.45%).

Treatments T₀, T₁, T₂, T₃, T₄, T₅ and T₆ were differ from one another considerably ($p < 0.05$).

Pardhi *et al.* (2014) reported that total sugars content of T₀, T₁, T₂ and T₃ was 14.43, 14.83, 15.51, 16.26 per cent, respectively and it differed significantly ($p < 0.05$) from each other.

Conclusion

The better quality lassi can be prepared by incorporating 6 per cent grape juice and 15 per cent sugar (overall acceptability score 8.47). The sensorily superior fresh lassi sample had average chemical composition 3.09 per cent fat, 3.26 per cent protein, 78.08 per cent moisture, 21.92 per cent total solids, 0.75 per cent ash, 0.85 per cent acidity, 4.34 per cent reducing sugars, 10.48 per cent non-reducing sugar, 14.82 per cent total sugars and 4.15 pH.

Reference

1. Avtade NB, Wankhede SA, Chaudhary DM. Effect of pomegranate juice on chemical quality of lassi. *Mysore J Agric Sci.* 2010;44(1):209-212.
2. Bagal S, Todkar S, Thalkar M. Utilization of papaya pulp for lassi preparation. *Adv Life Sci.* 2018;5(20):15-18.
3. Bendaali Y, Vaquero C, Gonzalez C, Morata A. Elaboration of an organic beverage based on grape juice with positive nutritional properties. *Food Sci Nutr.* 2022;10:1768-1779.
4. De S. *Outlines of Dairy Technology.* 29th ed. Delhi: Oxford Univ Press; 2004. p. 419.
5. Dhumal VS, Padghan PV, Shinde SP, Maske TA. Effect of pudina extract on physico-chemical properties of lassi with optimized level of pudina leaves. *J Pharmacogn Phytochem.* 2018;7(1):2763-2766.
6. Gaikwad J, Gubbawar SG, Kadu K, Bhondave A. Utilization of menthol (*Mentha arvensis*) juice for preparation of lassi from cow milk. *Int J Chem Stud.* 2018;6(4):2330-2332.
7. Ghule BK, Desale RJ, Gavhane MS, Khore MC. Preparation of strawberry lassi. *Res J Anim Husb Dairy Sci.* 2015;6(1):22-26.
8. Hapse PS. Preparation of chakka whey beverage using mango pulp [MSc thesis]. Rahuri: MPKV; 2004.
9. Hossain MN, Fakruddin M, Islam MN. Development of fruit dahi (yoghurt) fortified with grapes juice. 2012.
10. Jadhav DS, Chavan KD, Desale RJ. Influence of addition of orange (*Citrus sinensis*) juice on sensory and microbiological qualities of lassi during storage. *Asian J Dairy Food Res.* 2014;33(3):187-189.
11. Kedaree VC, Nalkar SD, Deshmukh MB. Preparation of lassi blended with kiwi (*Actinidia deliciosa*) pulp. *J Pharmacogn Phytochem.* 2021;10(2):768-771.
12. Kersh E, Hammad G, Donia MS, Farag MA. A comprehensive review on grape juice beverage in context to its processing and composition with future perspectives to maximize its value. *Food Bioprocess Technol.* 2023;16(1):1-23.
13. Khore NC. Preparation of strawberry lassi [MSc thesis]. Rahuri: MPKV; 2013. p. 41-2.
14. Matkar SP. Studies on preparation of papaya lassi with special reference to lactic acid bacteria [PhD thesis]. Rahuri: MPKV; 2010.
15. Navale MS. Utilization of orange (*Citrus reticulata*) juice for preparation of lassi [Doctoral dissertation]. Akola: Dr. Panjabrao Deshmukh Krishi Vidyapeeth; 2020.
16. Pardhi PS, Desale RJ, Mule PR, Ghule BK, Tambe DR, Gavhane MS. Studies on finger millet lassi. *Asian J Dairy Food Res.* 2014;33(4):255-8.
17. Prabhakar VC. Preparation of low-fat, mango-fortified bajara lassi [MSc thesis]. Rahuri: MPKV; 2018.
18. Roy JR, Sinthia ST, Roy JR, Abdul M, Sheikh M, Roy J, Mahomud MS. Effect of grape juice on the physiochemical and sensory properties of yogurt. *J Environ Sci Toxicol Food Technol.* 2019;13(9):94-100.
19. Singh J, Amin J. Role of microorganism in fermented milk. *Dairy Guide.* 1983;5(1):26-30.
20. Shiraishi M, Fujishima H, Chijiwa H. Evaluation of table grape genetic resources for sugar, organic acid, and amino acid composition of berries. *Euphytica.* 2010;174(1):1-13.
21. Tambe NN, Gubbawar SG, Mane NG, Mane RL, Raskar YM. Preparation of papaya (*Carica papaya* L.) pulp lassi from buffalo milk. *Trends Biosci.* 2017;10(36):7595-8.
22. Upadhyay S, Parimita T, Kumar P. Preparation of carrot lassi. *Pharma Innov J.* 2017;6(8):302-5.
23. Zare S, Lashkari H. Production and characterization of flavored dairy dessert containing grape juice concentrate. *Iran J Chem Chem Eng.* 2021;40(6):2028-41.